



Post-Operative Instructions for Implants

Please follow these instructions as closely as possible. They are designed to help you heal quickly and comfortably, with minimal side effects. If you experience any complications, please call our office and we will return your call as soon as possible.

For the remainder of the day of surgery:

- 1) Do not spit. Use a tissue to wipe your mouth as needed or swallow your saliva.
- 2) Do not use a drinking straw. Drink straight from cup.
- 3) Do not smoke.
- 4) Keep fingers and tongue away from the surgical area. Spitting, the use of a straw, smoking, and poking can dislodge the blood clot that is forming and may cause bleeding from the area. Also, smoking can increase the chances of an infection.

For Bleeding: Some minor bleeding is expected after implant surgery. It will usually subside quickly, and stop within an hour or two after surgery. A little oozing is normal and may persist for several hours. Upper implants may occasionally trigger some bleeding from the nose and this is normal and will stop quickly.

- 1) Keep gauze on the surgical area with some pressure for 30 to 45 minutes.
- 2) Remove the gauze after 30 to 45 minutes, and replace it with new piece of gauze if you are still bleeding. It is important to make sure the gauze is directly on the surgical site. Firm pressure for another hour should stop the bleeding.
- 3) If you find that this is not working after your third attempt, use a moistened tea bag instead of the gauze. The tannic acid in tea will help to form a clot and stop the bleeding.
- 4) If still cannot control the bleeding call the office.

For Swelling: Most patients will experience some swelling after surgery in the mouth. It may be mild or severe and is different for every patient. The swelling will get bigger for the first 24 to 48 hours before it starts to go away. It may last for several days to one week. Some bruising may also develop on the face.

- 1) Use an ice pack on the cheek or face next to the surgical site. Keep it on for 15 to 20 minutes, then off for 10 minutes, and repeat until you go to sleep. Start again the next day.

- 2) Sleep with your head elevated slightly, above the heart. This will keep swelling down.
- 3) On the third day, change to moist heat instead of ice packs. This will bring the swelling down quicker.

Eating: You may eat soft foods as soon as the anesthetic wears off. Do not to chew directly on the surgical site. You may resume a regular diet as soon as you feel up to it. Please stay well nourished and well hydrated; you will heal faster.

Brushing: You may brush your teeth, avoiding the surgical area, either tonight or tomorrow morning. Be gentle and do not spit or rinse forcefully. Start brushing the surgical area on the second day and be very gentle on the stitches.

Rinsing: You may start rinsing today, gently, with some warm salty water every few hours. Do not use alcohol-containing mouth rinses for a few days.

Medications: If you were prescribed any medications, take as directed. Call us if you experience severe nausea, diarrhea, or cannot swallow your pills.

- 1) Antibiotics: continue until the bottle is empty. Do not quit halfway.
- 2) Pain Medicine: continue as necessary. Remember that narcotics can make you drowsy so no driving, operating machinery, or alcoholic beverages while you are taking them.
- 3) Mouth rinse: Peridex or Periogard. Swish with half ounce for thirty seconds then spit it out, twice daily.

Pain Management:

1st level (mild-moderate) - If you have a prescription for Ibuprofen please take as instructed. Take (800mg) Ibuprofen every 6-8 hours. Take Tylenol every 4 hours. Alternate medicines.

2nd level (Severe) - If necessary, take the prescribed pain medication as directed. Continue taking Ibuprofen and substitute the prescribed pain medicine for the tylenol.

Example: (1) 800 mg Ibuprofen every 8 hours.
(1) 325 mg Tylenol every 4 hours.
(1) Prescribed pain pill every 4-6 hours substituting for Tylenol. *Do not exceed 4,000 mg of Acetaminophen or 3,200 mg of Ibuprofen per day